An Autopsy Study of Atherosclerotic Changes in Coronary Arteries at B.P. Koirala Institute of Health Sciences

Sudhir Raman Parajuli¹, Bishwonath Yadav¹, Prahlad Karki², Paricha Upadhyaya³, Shivendra Jha¹

¹Department of Forensic Medicine, B.P. Koirala Institute of Health Sciences, Dharan, Nepal

²Department of Internal Medicine, B.P. Koirala Institute of Health Sciences, Dharan, Nepal

³Department of Pathology, B.P. Koirala Institute of Health Sciences, Dharan, Nepal

CORRESPONDENCE

Dr. Sudhir Raman Parajuli,
Department of Forensic Medicine, Manipal
College of Medical Sciences, Pokhara,
Nepal
Email: drsudhirramanparajuli@yahoo.com

ARTICLE INFO

Article History

Submitted: 01 November, 2020

Accepted: 19 December, 2020

Published: 8 February, 2021

Source of support: None

Conflict of Interest: None

Copyright : ©The Author(S) 2020

This is an open access article under
the Creative Commons Attribution
license CC-BY 4.0

ABSTRACT

Introduction: Atherosclerosis is a giant killer and the incidence of atherosclerosis in coronary arteries is rapidly increasing. The study was designed to assess the atherosclerotic lesions in coronary artery and to correlate the risk factors related to prevalence of atherosclerosis.

Methods: Heart from 100 medico legal autopsy cases ranging between ages 15 to 35 years which came to BPKIHS Dharan were taken for this study and processed for coronary arteries using conventional technique. They were then studied, and grading was done based on Modified American Heart Association (AHA) classification of atherosclerosis.

Results: Intimal thickening was noted in more than 90% in all three coronary arteries followed by intimal xanthoma whereas intermediate lesion for atherosclerosis was not found. Age, gender, smoking and alcohol in relation to atherosclerosis were found to be of no significance.

Conclusion: The study highlights the impact of atherosclerotic lesions in the Eastern region of Nepal. Meticulous postmortem examination along with histopathological study is the best possible way to study atherosclerotic disease in humans and risk factors associated with it.

Keywords: Atherosclerosis; Autopsy; Coronary artery.

INTRODUCTION

Coronary atherosclerosis is the most frequent cause of ischemic heart disease. The composition and vulnerability of the atherosclerotic plaque determines the development of acute coronary syndromes.¹ Inflammation has a key role in the pathogenesis of atherosclerosis and it is supported by numerous factors such as modified LDL, hypertension, diabetes mellitus, free radicals and, in particular, by infectious agents such as *Chlamydia pneumoniae*.² Atherosclerosis is characterized by intimal lesions called atheromas, agents such as *Chlamydia* agents lesions called atheromas, atheromatous or fibro fatty plaques, which protrude into and obstruct vascular lumens and weaken the underlying media.³

Atherosclerosis can affect the kidneys either directly (e.g. renal artery stenosis) or as a frequent site of atheroembolic disease.⁴ Atherosclerosis is a giant killer and the incidence of atherosclerosis in coronary arteries is rapidly increasing. Myocardial infarction alone is responsible for 20% to 25%

of all deaths in United States.³ The incidence of CAD is less common, amongst Nigerian as compared to the other races. In India, it accounts for 85% of all the sudden deaths.⁵ This study is defined to assess the histomorphological atherosclerotic changes in coronary arteries autopsy by applying the modified AHA classification of atherosclerosis based on morphological descriptions to find out the age, gender and other risk factors related prevalence of atherosclerosis in Eastern region of Nepal.⁶

This specific classification is used since it offers better categorization of atherosclerotic lesions based on morphological descriptions compared to the earlier AHA classification, which was too rigid with descriptions of lesions (Table 1). Defining these intermediate lesions is of great importance as they are the precursor lesions of future advanced atherosclerosis.

METHODS

The present study is a hospital based cross sectional study carried out in dead bodies brought to BPKIHS mortuary from Eastern region of Nepal and was carried out from September 2014 to August 2015. Ethical clearance for the study was taken from Ethical Review Board, BPKIHS before the commencement of the study. Autopsy subject was identified, examined, and postmortem examination was done after obtaining requisition and inquest paper from the police in medico legal autopsies and after that cause of death were made. Historical data regarding age, gender, past medical illness, medications, familial and drug habits, habits such as alcohol consumption and smoking were noted from the deceased relatives and from the death certificate or police where available.

Heart and three coronary arteries: right (RCA), left anterior descending artery(LAD) and left circumflex arteries(LCA) were then examined grossly and each sectioned at 3-5 mm intervals to examine for any atherosclerotic plaques through microscopic assessment and grading was done according to modified AHA classification of atherosclerosis based on morphological descriptions for individual artery. Sample size of the study was 100. Data was collected and entered in MS EXCEL 2013 and analysis using SPSS 11.0 version.

Table 1: Modified AHA classification of atherosclerosis based on morphological description

Description	Thrombosis
Non atherosclerotic lesions	
Intimal thickening	Absent
Intimal xanthoma, or "fatty streak"	Luminal accumulation of foam cells without a necrotic core or fibrous cap. Based on animal and human data, such lesions usually regress.
Progressive atherosclerotic lesions	
Pathological intimal thickening	SMCs in a proteoglycan-rich matrix with areas of extra cellular lipid accumulation without necrosis
Erosion	Luminal thrombosis; plaque same as above
Fibrous cap atheroma	Well-formed necrotic core with an overlying fibrous cap
Erosion	Luminal thrombosis; plaque same as above; no communication of thrombus with necrotic core

Thin fibrous cap atheroma	The normal accumulation of Smooth Muscle Cells (SMCs) in the intima in the absence of lipid or macrophage foam cells	Absent
Plaque rupture	A thin fibrous cap infiltrated by macrophages and lymphocytes with rare SMCs and an underlying necrotic core	Absent; may contain intra-plaque hemorrhage/fibrin
Calcified nodule	Fibro atheroma with cap disruption; luminal thrombus communicates with the underlying necrotic core	Thrombus usually occlusive

RESULTS

A total of 406 autopsies were performed in the department of forensic medicine in one year from august 2014 to august 2015. Of those 100 cases were taken of age groups 15 to 35 years irrespective of the cause of death. The mean age was 24.81 years with standard deviation of 5.88. Majority of the population were literate being 25% of the total population.

Table 2: Coronary artery atherosclerotic lesions

Lesions	RCA	LAD	LCA
Intimal thickening	88	90	96
Intimal xanthoma	11	10	4
Progressive intimal thickening	1	0	0
Total	100	100	100

Major cause of death in this age group was hanging followed by road traffic accident and poisoning. Majority of the population were homemaker followed by self-employed and student. Unemployed and farmer share equal percentage of population of 9%. 13% of the population were unemployed. About half of the population group(52%) were smokers. More than half of the population(56%) consumed alcohol. (Figure 1) Intimal thickening was noted in most of the individuals followed by intimal xanthoma (Table 2).

Figure 1: Cause of death among study group

Fig : 2a

Fig : 2b

Fig : 2c

Figure 2

Microphotograph showing coronary artery Intimal thickening(2a), intimal xanthoma(2b) and Progressive intimal thickening (2c) using Hematoxylin and Eosin stain.

In this study, we tried to show association of atherosclerotic lesions with various other socio-demographic and behavioral predictors (Table 2a, 2b and 2c).

Table 2a: Association of LAD with age and gender

Characteristics	Category	LAD		p-value
		Intimal thickening	Intimal xanthoma	
Age	15-25		3(3.2%)	0.090
	26-35	335(83.3%)	7(16.7%)	
Gender	Male	355(94.8%)	6(10.7%)	1.000
	Female	40(90.9%)	4(9.1%)	

Table 2b: Association of LCA with age and gender

Characteristics	Category	LCA		p-value
		Intimal thickening	Intimal xanthoma	
Age	15-25	56(96.6%)	2(3.4%)	1.000
	26-35	40(95.2%)	2(4.8%)	
Gender	Male	53(94.6%)	3(5.4%)	0.628
	Female	43(97.7%)	1(2.3%)	

In this study no significant association was seen between age and atherosclerotic lesions in RCA, LAD and LCA. Similarly, although males were seen having more intimal xanthomatous lesions in LAD and LCA, this was however not significant.

Table 2c: Association of RCA with age and gender

Characteristics	Category	RCA		p-value
		Intimal thickening	Intimal xanthoma	
Age	15-25	52(89.7%)	6(10.3%)	0.552
	26-35	36(85.7%)	6(14.3%)	
Gender	Male	50(89.3%)	6(10.7%)	0.760
	Female	38(86.4%)	6(13.6%)	

Among the alcohol consumers, intimal xanthoma was seen in 7.1% of the cases, and this was not found to be statistically significant ($p=0.328$) in LAD. Smoking status was also not found to be associated with atherosclerotic lesions. (Table 3a, 3b, 3c)

Table 3a: Association of LAD with Alcohol consumption

Characteristics	Category	LAD		p-value
		Intimal thickening	Intimal xanthoma	
Alcohol	Yes	52(92.9%)	4(7.1%)	0.328
	No	38(86.4%)	6(13.6%)	
Smoking	Yes	48(92.3%)	4(7.1%)	0.514
	No	42(87.5%)	6(13.6%)	

Table 3b: Association of LCA with Alcohol consumption and smoking

Characteristics	Category	LCA		p-value
		Intimal thickening	Intimal xanthoma	
Alcohol	Yes	55(98.2%)	1(1.8%)	0.317
	No	41(93.2%)	3(6.8%)	
Smoking	Yes	51(98.1%)	1(1.9%)	0.348
	No	45(93.8%)	3(6.2%)	

Table 3c: Association of LAC with Alcohol consumption

Characteristics	Category	LAC		p-value
		Intimal thickening	Intimal xanthoma/ progressive intimal thickening	
Alcohol	Yes	50(89.3%)	6(10.7%)	0.223
	No	38(86.4%)	6(13.6%)	
Smoking	Yes	48(92.3%)	4(7.7%)	0.348
	No	40(83.3%)	8(16.7%)	

DISCUSSION

Coronary artery disease (CAD) is associated with the number of risk factors causing coronary atherosclerosis. Coronary artery stenosis is mostly caused by coronary

atherosclerosis.⁷ Coronary Heart Disease incidence is decreasing in Western Europe, the United States and in Australia but is steeply increasing in Asia, Central and Eastern Europe, and Africa.⁸ In past few years, Eastern region of Nepal has undergone a rapid change in lifestyle, habit modification, stress, strain. In the present study sufficient effort has been made to study the effects of these habits on atherosclerosis. Atherosclerosis is a multi-factorial disease affected by various factors, such as smoking, alcohol consumption, presence of stress and strain in life, personality characteristics, etc. Many studies have been undertaken in the past to study association of various factors leading to atherosclerosis, ultimately leading to CAD.^{5,9} In a study done by Adhikari CM et al. in a capital city heart center 70% patients were smokers who were brought for coronary heart disease in age group up to 40 years.¹⁰ Study done in Central Nepal by Koju R et al. showed that among the established cardiovascular risk factors, gender, diabetes mellitus and smoking are significantly associated with coronary stenosis among CAD patients which is in contrast to our study.⁷ In our study, non-atherosclerotic lesions were seen in almost all cases except in one case where the right coronary artery showed atherosclerotic lesion. A similar study using same classification conducted by Viral M. Bhanvadia et al. showed non-atheromatous lesions of 40% and atherosclerotic lesion of 60%. Our study showed a higher degree of involvement of atherosclerosis in LAD (42%) as compared to LCA (40%) and RCA (39%).¹¹

This study also showed contrast finding to that of Sudha et al where atherosclerotic lesions in LAD contributed (47%). Also, study carried out by Thej et al. and Yazdi et al. in Iran, showed more involvement of LAD (60%) in atherosclerosis in comparison to RCA (50%), and LCA (42.5%) respectively.^{12,6,13} Garg M et al. in his study concluded that atheroma have been seen above the age of 15 years, but significant atheroma appeared after third decade onwards and thereafter there is a gradual increase in both its frequency and severity from second to sixth decade. Commonest type of atherosclerosis was grade 3 described as intermediate lesion or pre-atheroma. Our study showed no atherosclerotic lesions in almost all subjects. This finding is surprisingly opposite to that of the studies conducted in different parts of the world. This could be attributed to the fact that in our study the subjects were free from diabetes, hypertension, and other cardiac illness despite being smokers and alcoholics.

In recent years, Magnetic Resonance Imaging (MRI) has emerged as a very promising technique for studying atherosclerotic diseases in humans, and various studies have proved their association with histopathological findings. Study done by Everis P. S et al. demonstrated good

agreement between coronary artery morphometrical measurements obtained by high resolution MRI and histopathology in an ex-vivo experimental model.¹⁵ In a country like Nepal where facility of MRI is limited and costly, investigation with these modalities are beyond the capacity of people in both living and dead. In Nepal autopsy-based research pertaining to atherosclerosis and its associated risk factors is lacking. Therefore, autopsy and histopathological based studies in this topic covering a larger sample could be beneficial in confirming the association between atherosclerosis and its risk factors.

CONCLUSION

The study showed unexpectedly lower prevalence of atherosclerosis in younger age group with no significant association between atherosclerosis and its risk factors. Further, study with large sample size is required to better elucidate the association between the two. Histopathological studies provide the most accurate clues to a better understanding of human coronary artery disease. Hence, meticulous postmortem examination along with histopathological study is the best possible way to study atherosclerotic disease in humans and risk factors associated with it.

REFERENCE

1. Amel E, Prathiba D, Kumar S. Morphological and morphometric analysis of coronary atherosclerosis--an autopsy study. *Indian J Pathol Microbiol.* 2006;49(2):239–42.
2. Gaudio E, Carpino G, Grassi M, Musca A. Morphological aspects of atherosclerosis lesion: past and present. *Clin Ter.* 157(2):135–42.
3. Kumar, Abbas, Fausto. Robbins and Cotran Pathological Basis of Disease. 7th ed. 516 p.
4. Kasper et al. Harrison's Principles of Internal Medicine. 16th ed. 1425 p.
5. Singh H, Oberoi S, Gorea R, Bal M. Atherosclerosis in coronaries in Malwa Region of Punjab. *JIAFM.* 2005;27(4):236–9.
6. Thej MJ, Kalyani R, Kiran J. Atherosclerosis in coronary artery and aorta in a semi-urban population by applying modified American Heart Association classification of atherosclerosis: An autopsy study. *J Cardiovasc Dis Res.* 2012 Oct;3(4):265–71.
7. Koju R, Humagain S, Khanal K. Association of cardiovascular risk factors and coronary artery lesion among coronary

- artery disease patients. Kathmandu Univ Med J. 12(46):137-40.
7. Koju R, Humagain S, Khanal K. Association of cardiovascular risk factors and coronary artery lesion among coronary artery disease patients. Kathmandu Univ Med J. 12(46):137-40.
 8. Assmann G, Cullen P, Jossa F, Lewis B MM. Coronary heart disease: Reducing the risk: The scientific background to primary and secondary prevention of coronary heart disease. A worldwide view. International task force for the prevention of coronary heart disease. Arter Thromb Vasc Biol. 1999;19:1819-24.
 9. Abraham Joseph, MD, FACC, Douglas Ackerman, MD, J. David Talley, MD, FACC, John Johnstone, MD, FACC, Joel Kupersmith, MD F. Manifestations of Coronary Atherosclerosis in Young Trauma Victims-An Autopsy Study. JACC. 22(2):459-67
 10. Adhikari, C., Rajbhandari, R., Limbu, Y., Malla, R., Sharma, R., Rauniyar, B., Rajbhandari, S., Baidya, S., Sharma, D., Maskey, A., & KC, M. (2013). A study on major cardiovascular risk factors in Acute Coronary Syndrome (ACS) patient 40 years and below admitted in CCU of Shahid Gangalal National Heart Center. Nepalese Heart Journal, 7(1), 20-4.
 11. Viral M. Bhanvadia et al. Study of Coronary Atherosclerosis by Modified American Heart Association Classification of Atherosclerosis-An Autopsy Study. J Clin Diagnostic Res. 7(11):2494-7.
 12. Sudha M L, Sundaram S, Purushothaman K R, Kumar P S PD. Coronary atherosclerosis in sudden cardiac death: An autopsy study. Indian J Pathol Microbiol. 2009;52:486-9.
 13. Yazdi A, Rezaei A, Azari J, Hezazi A, Shakeri M, Shahri M. prevalence of atherosclerotic plaques in autopsy cases with non cardiac death. Iran J Pathol. 2009;4:101-4.
 14. Garg M, Aggarwal AD, Kataria SP. Original Research Paper Coronary Atherosclerosis and Myocardial Infarction An Autopsy Study. 2011;33(1):971-4.
 15. Gomes EPSG, Rochitte CE, Azevedo CF, Lemos PA, Gutierrez PS, César LAM. Ex-vivo Assessment of Coronary Artery Atherosclerosis by Magnetic Resonance Imaging: Correlation with Histopathology. Open Cardiovasc Med J. 2014;8:26-34.